arXiv:math/0701036v2 [math.AG] 3 Jan 2007

A SIMPLE APPLICATION OF ENRIQUES-SEVERI-ZARISKI-SERRE TO THE LOCAL FUNDAMENTAL GROUP SCHEME

V. B. MEHTA

Abstract. We prove that the Local Fundamental Group Scheme satisfies the Lefschetz- Bott theorems in characteristic p. The proofs are standard applications of the E-S-Z-S vanishing theorems and are routine.

1. Introduction

The Fundamental Group Scheme was introduced by Madhav Nori in [N1,N2]. In order to prove the conjectures made in [loc.cit], S.Subramanian and the present author had introduced the "Local Fundamental Group Scheme ", denoted by $\pi^{loc}(X)$[MS1], which is the infinitesimal part of $\pi(X)$,[MS2]. One may ask whether the Lefschetz-Bott theorems hold for $\pi^{loc}(X)$. In other words, let X be a smooth projective variety over an algebraically closed filed of characteristic p,and let H be a very ample line bundle on X. The question is if $\pi^{loc}(Y) \to \pi^{loc}(X)$ is a *surjection* if dim $X \geq 2$ and degree $Y \geq n_0$, where n_0 is an integer depending *only* on X. Similiarly , if dim $X \geq 3$,the question is if $\pi^{loc}(Y) \to \pi^{loc}(X)$ is an *isomorphism* if deg $Y \geq n_1$, where n_1 is an integer depending *only* on X.

In the present note, we give a positive answer to both these questions. The methods only involve applying the lemma of Enriques-Severi-Zariski-Serre (E-S-Z-S) over and over again. Hence most of the proofs are only sketched, and some are completely omitted. The notation is the same as in [MS2],which is briefly recalled.

2. The first theorem

Theorem 2.1. *Let X be a non-singular projective variety over an algebraically closed field of char p,of dimension ≥ 2. Let H be a very ample line bundle on X. Then there exists an integer n_0, depending only on X, such that for any smooth $Y \in |nH|$, of deg $n \geq n_0$, the canonical map : $\pi^{loc}Y \to \pi^{loc}(X)$ is a surjection.*

Before we begin the proof, we recall some facts from [MS2]. Let $F : X \to X$ be the Frobenius map. For any integer $t \geq 1$, denote by C_t the category of all $V \in Vect(X)$ such that $F^{t*}(V)$ is the *trivial* vector bundle on X.Fix a base point $x_0 \in X$, Consider the functor $S : C_t \to Vect(X)$ given by $V \to V_{x_0}$ It is seen that C_t , with the fibre functor S, is a Tannaka category [MS1,MS2]. denote the corresponding Tannaka group by G_t. It is also easily seen that

$$\pi^{loc}(X) \simeq \lim_{\leftarrow t} G_t$$

1991 *Mathematics Subject Classification.* 14F20,14L15.

, where $\pi^{loc}(X)$ is the Tannaka group associated to the category $(V \in Vect(X)\ F^t(V)$ is trivial for some t.

Now let $t = 1$, and consider $G_1(Y)$ and $G_1(X)$. Let $V \in C_1(X)$. with V stable.V corresponds to a principal H bundle $E \to X$,where E is *reduced* in the sense of Nori [N1,page 87,Prop.3], or just N-reduced. To prove that $G_1(Y) \to G_1(X)$ is a surjection , it is enough to prove that V/Y is *stable*, or better still that E/Y is *N-reduced.*

Lemma 2.2. *If X and E are as above , then there exists an integer n_0, depending only on X ,such that for all $n \geq n_0$, and for all smooth $Y \in |nH|$, E/Y is N-reduced.*

Proof. Let $f : E \to X$ be given.Then $f_*\mathcal{O}_E$ belongs to C_1. Denote it by W for simplicity. Now consider

$$0 \to \mathcal{O}_X \to F_*\mathcal{O}_X \to B^1 \to 0 \tag{1}$$

, Tensor (1) with $W(n)$. So there is an n_0 such that for $n \geq n_0$, we have $Hom(W(n), B^1) = 0$. So the canonical map $H^1(W(n) \to H^1(F^*(W(n))$ is injective. But the last space is just $H^1(\mathcal{O}_X(np))$ which is 0 as soon as $n \geq n_1$, independent of $V \in C_1(X)$. Hence $H^0(Y, W/Y) = 1$, which proves that E restricted to Y is also N-reduced.

Now assume $t \geq 2$. We shall assume in fact that $t = 2$, because an almost identical proof works for $t \geq 3$.Consider the map $F^2 : X \to X$. Let B_2^1 be the cokernel.We have the exact sequence

$$0 \to B^1 \to B_2^1 \to F_*B_1 \to 0 \tag{2}$$

Let $W \in C_2(X)$ and tensor (2) with $W(n)$. One sees immediately that if $H^1(W(n)\otimes B^1) = 0$ for all $n \geq n_0$, then also $H^1((W(n)\otimes B_2^1) = 0$ for all $n \geq n_0$, for the *same* integer n_0.So if E is N-reduced on X, then E remains N-reduced on Y, if deg $Y \geq n_0$, for the same integer n_0, which worked for $C_1(X)$.The proof for bigger t goes the same way , by taking the cokernel of F^t, where $t \geq 3$. This concludes the proof of Theorem 2.1. □

3. THE SECOND THEOREM

Theorem 3.1. *Let X be smooth and projective of dim ≥ 3. Then there exists an integer n_0, depending only on X ,such that for any smooth $Y \in |nH|, n \geq n_0$, the canonical map $\pi^{loc}(Y) \to \pi^{loc}(X)$ is an isomorphism.*

Proof. We assume that dim $X \geq 3$, and we pick an arbitrary smooth $Y \in |nH|$.We first consider a $W \in C_1(Y)$ and show that it lifts uniquely to $V \in C_1(X)$. Such a W is trivialized by the Frobenius,so there exists M, an $r \times r$ matrix of 1- forms on Y,which gives an integrable connection ∇ on $\mathcal{O}_Y^r$, with p- curvature 0. Consider

$$0 \to \frac{I}{I^2} \to \Omega_X^1/Y \to \Omega_Y^1 \to 0 \tag{1}$$

and

$$0 \to \Omega_X^1(-n) \to \Omega_X^1 \to \Omega_X^1/Y \to 0 \tag{2}$$

Here I is the idealsheaf of $Y, I = \mathcal{O}_X(-n)$. We get an integer n_0, such that for all $n \geq n_0, H^0(X, \Omega^1_X) \to H^0(Y, \Omega^1_Y)$ is an isomorphism. So M lifts to M_1, a $r \times r$ matrix of 1 forms on X.Now ∇ has p-curvature 0 on Y.We now show that ∇_1 defined by M_1 has p-curvature 0 on X. The curavture of ∇ is an element of $H^0(End\mathcal{O}^r_Y \otimes \Omega^2_Y)$, Again by E-S-Z-S, the curvature of ∇_1 is 0 if deg $Y \geq n_1$, for some integer n_1, depending only on X. The p-curvature of ∇_1 is an element of $H^0(F^*\Omega^1_X \otimes End\mathcal{O}^r_X)$, which again vanishes if deg $Y \geq n_2$. So any element $W \in C_1(Y)$ lifts uniquely to an element $V \in C_1(X)$.

Lemma 3.2. *With X as above, then there exists an integer n_3 ,depending only on X, such that for any $V \in C_t(X)$, any t, we have $H^1(\Omega^1_X(-n) \otimes V) = 0$, for all $n \geq n_3$.*

Proof. Let T_X be the tangent bundle of X. For some s , depending only on X, $T_X(s)$ is generated by global sections. So we have

$$0 \to S^* \to \mathcal{O}^N_X \to T_X(s) \to 0.$$

Dualizing,we get

$$0 \to \Omega^1(-s) \to \mathcal{O}^N_X \to S \to 0 \tag{1}$$

.Tensor (1) with $V(-t)$,to get

$$0 \to V(-t) \otimes \Omega^1(-s) \to V(-t) \otimes \mathcal{O}^N_X \to V(-t) \otimes S \to 0 \tag{2}$$

. Applying F to (2) we get

$$0 \to [V(-t) \otimes \Omega^1(-s)]^p \to [V(-t) \otimes \mathcal{O}^N_X]^p \to [V(-t) \otimes S]^p \to 0 \tag{3}$$

. It is clear that $H^0(V(-t) \otimes S) = 0$ for $t >> 0$, *independent* of V. Now look at

$$H^1(V(-t) \otimes \Omega^1(-s)) \to H^1(V(-t) \otimes \mathcal{O}^N_X) \to H^1(V(-t) \otimes \mathcal{O}^N_X)^p \to 0 \tag{4}$$

,where the right arrow is the map induced by F.But one knows that $H^1(V(-t) \otimes \mathcal{O}^N_X)^p$ vanishes for $t >> 0$, independently of $V \in C_n(X)$ for any n. Therefore $H^1(V(-t) \otimes \Omega^1)$ vanishes for all $t >> n$, for an integer n *independent* of X.This concludes the proof of Lemma 3.1 ☐

Proceeding with the proof of Theorem 3.1 by the discussion preceeding the Lemma 3.1,it follows that if $W \in C_1(Y)$ lifts to V on X,such that $V \in C_1(X)$ and W has a p-flat connection ,then V also has a connection.The integrability of this connection on V is also defined by the vanishing of a tensor in $H^0(EndV \otimes \Omega^2_X)$. It is again clear that this tensor vanishes if its restriction to Y vanishes, for Y of degree say n_4 ,again independent of $V \in C_t(X)$, for all t. Again the p-flatness of this connection on V also vanishes ,if the corresponding p-curvature vanishes for V restricted to Y, for all smooth $Y \in |nH|$, for n independent of V. This means that on X, if V restricts to W on Y, and W has a p-flat connection, then V also has a p-flat connection. So if W on Y descends to W_1 , then V also descends under F to V_1, and that V_1 restricts to W_1 on Y. Clearly ,this continues to hold for $V \in C_t(X)$, restricting to $W \in C_t(Y)$, any t,for Y of a fixed degree ,depending *only* on X. Hence the canonical map of Tannaka Categories : $C_t(X) \to C_t(Y)$, given by restriction from X to Y, induces an *isomorphism*:

$$G_t(Y) \to G_t(X),$$

for all t.But

$$\pi^{loc}(X) \simeq \lim_{\leftarrow t} G_t(X)$$

and similiarly for Y. This finishes the proof of Theorem 3.1 $\square$

Remark 3.3. It would be of great interest to prove the above theorems when X is only assumed to be *normal.*

Remark 3.4. It is also interesting to determine if the category of F-trivial vector bundles ,on a smooth X,is m- *regular*. This is the case if dim $X \leq 3$.

REFERENCES

[MS1] V. B. Mehta, S.Subramanian *On the Fundamental group scheme*, Invent. Math. **148**,(2002),143-150.

[MS2] V.B. Mehta, S.Subramanian *Some Remarks on The Local Fundamental Group Scheme* submitted to the Proc. of the Ind. Acad. of Sciences, (Math Sci).

[N1] M.V.Nori *The Fundamental Group Scheme*,Proc. of the Ind. Acad. of Sci.(math Sci.),**91**,1982.

[N2] M.V.Nori *Representations of the Fundamental Group*,Compositio Math.**33**, 1976.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI-400005, INDIA

E-mail address: `vikram@math.tifr.res.in`